



Task Trol

Stjepan recently received his bachelor's degree in mathematics from the University of Zagreb. Naturally, his parents are very proud and have decided to give him all positive integers not greater than 2^{60} as a gift. To keep them safe, he quickly stored all of those numbers in an array A , such that $A_i = i$.

His jealous friend Marin decided to prank him by repeatedly replacing each element of A with the sum of its digits until all elements of A consisted of a single digit. For example, the initial value of 197th element of A was 197. Marin first changed that value to $1 + 9 + 7 = 17$ and then changed its value again to $1 + 7 = 8$.

Stjepan is devastated and begs Marin to return his array to its initial state. Unfortunately, Marin won't do that until Stjepan correctly answers his Q queries: "What is the sum of numbers from l -th to r -th element of A ?"

Help Stjepan answer those queries!

Input

The first line contains an integer Q ($1 \leq Q \leq 100$) from the task description.

The next Q lines contain two integers l_i i r_i ($1 \leq l_i \leq r_i \leq 2^{60}$), the parameters of Marin's i -th query.

Output

Output the answers to each of Marin's Q queries. Each answer should be printed in a separate line and their order should match the order of the queries as they are given in the input.

Scoring

In test cases worth a total of 10 points, for each query will hold $1 \leq l_i \leq r_i \leq 9$.

In test cases worth a total of 30 points, for each query will hold $r_i - l_i \leq 1000$.

Examples

input

1
1 5

output

15

input

2
9 13
44 45

output

19
17

input

1
1998 2018

output

102

Clarification of the second example:

1st query $\rightarrow A_9 = 9, A_{10} = 1 + 0 = 1, A_{11} = 1 + 1 = 2, A_{12} = 1 + 2 = 3, A_{13} = 1 + 3 = 4.$

$A_9 + A_{10} + A_{11} + A_{12} + A_{13} = 9 + 1 + 2 + 3 + 4 = 19.$

2nd query $\rightarrow A_{44} = 4 + 4 = 8, A_{45} = 4 + 5 = 9. A_{44} + A_{45} = 8 + 9 = 17.$

